

PHYSICS 112

Homework 5

Due in class, Tuesday February 14

Remember MIDERM:

In class on Thursday 10:00–11:10am. This will be followed by a short break and lecture.

1. Symmetry of filled and vacant orbitals

Consider the Fermi-Dirac distribution $f(\epsilon)$. Let $\epsilon = \mu + \delta$, so δ is the difference between the energy and the chemical potential. Show that

$$f(\mu + \delta) = 1 - f(\mu - \delta),$$

so the probability that an orbital ϵ above the chemical potential is occupied, is equal to the probability that an orbital δ below the chemical potential is empty. A vacant orbital is sometimes known as a **hole**.

2. Entropy of Mixing

Suppose that a system of N atoms of type A is placed in diffusive contact with a system of N atoms of type B at the same temperature and volume. Show that after diffusive equilibrium is reached, the entropy has increased by $2N \log 2$. This increase in entropy is called the *entropy of mixing*.

If the atoms are identical ($A \equiv B$), show that there is no increase in entropy when diffusive contact is established. This difference in results is called the *Gibbs paradox*.

3. Time for a large fluctuation

We ask how long it will take for a large fluctuation to occur in a macroscopic system. Consider a gas of ${}^4\text{He}$ atoms in a container of volume 0.1 liter at 300 K and a pressure of 1 atm. We ask how long it will take before the atoms are in a configuration in which they are all in one half of the container.

- (a) Estimate the number of states accessible to this system in this initial condition.

Note: For this part, and other parts of this question which involve *huge* numbers (i.e. numbers of order e^N where $N \sim 10^{21}$), you should express your answer in the form 10^x , with the appropriate value for x . This gives an intuitive appreciation of the result since x is the number of digits in the answer.

- (b) The gas is compressed isothermally to a volume of 0.05 liters (half of the initial volume). Estimate the number of states accessible now.
- (c) For the system in the 0.1 liter container estimate the value of the ratio

$$\frac{\text{number of states for which all atoms are in one-half of the volume}}{\text{number of states for which the atoms are anywhere in the volume}}.$$

- (d) If the collision rate of an atom is $\approx 10^{10} \text{ s}^{-1}$, what is the total number of collisions of all atoms in the system in a year? Use this as the frequency with which the state of the system changes.
- (e) Estimate the number of years you would have to wait for all atoms to be in one half of the system, starting from the equilibrium configuration.

Note: As stated in part (a), express your answer in the form 10^x and give the value of x .

4. **Gas of atoms with an internal degree of freedom**

Consider a classical ideal monatomic gas, where each atom has two internal energy states, one an energy Δ above the other. There are N atoms in volume V at temperature T . Find the

- (a) chemical potential,
- (b) free energy,
- (c) entropy,
- (d) pressure, and
- (e) heat capacity at constant pressure.