

PHYSICS 250
Homework 7
Due in class, Monday November 26.

1. Consider the following first order differential equation:

$$\frac{dx}{dt} + \gamma x(t) = F(t),$$

with $\gamma > 0$. In the absence of the “forcing term”, $F(t)$, the (homogeneous) equation, $x'(t) + \gamma x(t) = 0$, has solution

$$x(t) = Ae^{-\gamma t},$$

where A is a constant. As discussed in class, the solution of the inhomogeneous equation (i.e. including the forcing term) can be written as

$$x(t) = \int_{-\infty}^{\infty} G(t-t')F(t')dt',$$

where the Green function $G(t)$ satisfies

$$\frac{dG}{dt} + \gamma G(t) = \delta(t).$$

We assume the boundary condition that x is zero before the “force” $F(t)$ starts to act and so $G(t) = 0$ for $t < 0$.

- (a) Determine $G(t)$ either by working in the time domain, and determining the value of the constant A above needed to get the correct discontinuity at $t = 0$ due to the delta function, or use Fourier transforms (which automatically puts in the boundary condition that the solution vanishes at $t = \pm\infty$).
- (b) Denoting the Fourier transform of $G(t)$ by $\chi(\omega)$, determine $\chi'(\omega)$ and $\chi''(\omega)$, the real and imaginary parts, and verify, by doing the integrals, that they are related by Kramers-Kronig relations

$$\begin{aligned}\chi''(\omega) &= \frac{1}{\pi} \mathcal{P} \int_{-\infty}^{\infty} \frac{\chi'(\omega')}{\omega' - \omega} d\omega', \\ \chi'(\omega) &= -\frac{1}{\pi} \mathcal{P} \int_{-\infty}^{\infty} \frac{\chi''(\omega')}{\omega' - \omega} d\omega'.\end{aligned}$$

Note: Remember that the Kramers-Kronig relations are simply the result of causality, $G(t) = 0$ for $t < 0$) (and the fact that $\chi(\omega) \rightarrow 0$ for $|\omega| \rightarrow 0$).

2. Kelly, Qu. 7, Ch. 6.

Note: For part (c), you should find that

$$\begin{aligned}\bar{x}(t) &= \frac{\hbar k_0}{m} t, \\ \sigma^2(t) &= \sigma_0^2 + \left(\frac{\hbar t}{m \sigma_0} \right)^2.\end{aligned}$$

Note that at long times $\sigma(t) \propto t$. Physically, the reason for this is that different pieces of the wavefunction propagate with different velocities.

3. Kelly, Qu. 8, Ch. 6.

4. Kelly, Qu. 11, Ch. 6.

Note: Certain materials have regions of frequency where the refractive index, $n(\omega)$, is less than unity, implying a phase velocity, $v_p = c/n(\omega)$, greater than c , the speed of light in vacuum. The point of this question is for you to show that, nonetheless, no signal can propagate faster than c , in agreement with special relativity.

5. Kelly, Qu. 12, Ch. 6.

Note: For part (d) you are asked to *verify* that the given solution satisfies the equation provided certain relations between the parameters are satisfied. You may use *Mathematica* to help you with this.

6. Kelly, Qu. 13, Ch. 6.

Note: For part (c), you are only asked to *verify* the given solution, not to derive it “from first principles”. You may use *Mathematica* to help you, if you wish.