

Some recent developments in Spin Glasses

A. P. Young

Department of Physics, University of California, Santa Cruz, California 95064

Abstract. I give some experimental and theoretical background to spin glasses, and then discuss the nature of the phase transition in spin glasses with *vector* spins. Results of Monte Carlo simulations of the Heisenberg spin glass model in three dimensions are presented. A finite size scaling analysis of the correlation length of the spins and chiralities shows that there is a single, finite-temperature transition at which both spins and chiralities order.

Keywords. spin glass, Monte Carlo, phase transition

PACS Nos 75.50.Lk, 75.40.Mg, 05.50.+q

1. Introduction

For a theorist such as myself, a spin glass is a system with disorder and frustration. A simple but useful example is shown in Figure 1 in which there is a single square of Ising spins (which can only point up or down). The “+” or “-” on the bonds indicates a ferromagnetic or antiferromagnetic interaction respectively. In this example, with one negative bond, it is impossible to minimize the energy of all the bonds so there is competition or “frustration”. Clearly, for a large system, it is non-trivial to find even the ground state of a system with frustration and disorder.

The simplest model which has the basic ingredients of disorder and frustration is the the Edwards-Anderson [1] (EA) model,




Figure 1. A toy model which shows frustration. If the interaction on the bond is a “+”, the spins want to be parallel and if it is a “-” they want to be antiparallel. Clearly all these conditions can not be met so there is competition or “frustration”.

$$\mathcal{H} = - \sum_{\langle i,j \rangle} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j, \quad (1)$$

in which the spins \mathbf{S}_i lie on the sites of a regular lattice, and the interactions J_{ij} , which we take to be between nearest neighbors only, are independent random variables with mean and standard deviation given by

$$[J_{ij}]_{av} = 0; \quad [J_{ij}^2]_{av}^{1/2} = J (= 1). \quad (2)$$

A zero mean is chosen to avoid any bias towards ferromagnetism or antiferromagnetism, and it is convenient, in the simulations, to take a Gaussian distribution for the J_{ij} . The \mathbf{S}_i are of unit length and have m -components:

$$\begin{aligned} m = 1 & \quad (\text{Ising}) \\ m = 2 & \quad (\text{XY}) \\ m = 3 & \quad (\text{Heisenberg}). \end{aligned} \quad (3)$$

Experimentally, there are different types of spin glasses:

- Metals:
Diluted magnetic atoms, e.g. Mn, in a non-magnetic metal such as Cu, interact with the RKKY interaction,

$$J_{ij} \sim \frac{\cos(2k_F R_{ij})}{R_{ij}^3}, \quad (4)$$

where k_F is the Fermi wavevector. We see that J_{ij} is random in magnitude and *sign*, so there is frustration. Note that Mn is an S-state ion and so has little anisotropy. It should therefore correspond to a *Heisenberg* spin glass.

- Insulators:
An example is $\text{Fe}_{0.5}\text{Mn}_{0.5}\text{TiO}_3$, which comprises hexagonal layers. The spins align perpendicular to layers (hence it is *Ising*-like). Within a layer the spins in pure FeTiO_3 are ferromagnetically coupled while spins in pure MnTiO_3 are antiferromagnetically coupled. Hence the mixture gives an *Ising spin glass with short range interactions*.
- Other systems where spin glass ideas have proved useful are:
 - Protein folding
 - Optimization problems in computer science
 - Polymer glasses, foams ...

After considerable experimental and theoretical work, it became clear that a spin glass has a sharp thermodynamic phase transition at temperature $T = T_{SG}$, such that for $T < T_{SG}$ the spins freeze in some random-looking orientation. As $T \rightarrow T_{SG}^+$, the spin glass correlation length ξ_{SG} , which we will discuss in detail below, diverges. Here we just note that the defining feature of the correlation length is that the *magnitude* of the correlation




Figure 2. Results for the non-linear susceptibility of 1% Mn in Cu from Omari et al. [2]. The quantity a_3 is the non-linear susceptibility in dimensionless units.

function $\langle S_i S_j \rangle$ becomes significant for $R_{ij} < \xi_{SG}$, though the *sign* is random. A quantity which diverges, therefore, is the *spin glass susceptibility*

$$\chi_{SG} = \frac{1}{N} \sum_{\langle i,j \rangle} [\langle \mathbf{S}_i \cdot \mathbf{S}_j \rangle^2]_{av}, \quad (5)$$

(notice the square) which is accessible in simulations. It is also essentially the same as the *non-linear susceptibility*, χ_{nl} , which can be measured experimentally and is defined by the coefficient of h^3 in the expansion of the magnetization m ,

$$m = \chi h - \chi_{nl} h^3 + \dots, \quad (6)$$

where h is the magnetic field. We expect that χ_{nl} diverges at T_{SG} like

$$\chi_{nl} \sim (T - T_{SG})^{-\gamma} \quad (7)$$

where γ is a critical exponent.

This divergent behavior has been seen in many experiments. Fig. 2 shows the results of Omari et al. [2] on 1% Mn in Cu. They define $m = a_1 h - a_3 c_3 h^3 + a_5 c_5 h^5$ and choose units (and constants $c_3 = 1/15, c_5 = 2/305$) such that $a_i = 1$ for independent Mn spins. It follows that a_3 is χ_{nl} in dimensionless units. We see that χ_{nl} becomes very large, ($> 10^3$), and presumably diverges. A fit gives $\gamma = 3.25$.

At low temperatures, the dynamics of spin glasses becomes very slow, such that below T_{SG} the system is never fully in equilibrium. This is because the “energy landscape” becomes very complicated with many “valleys” separated by “barriers”. The (free) energies

of the valleys can be very similar and yet the spin configurations rather different. Hence there are large-scale, low-energy excitations in spin glasses.

This non-equilibrium behavior has been extensively studied in recent years. Of particular note has been the study of “aging” in spin glasses, pioneered by the Uppsala group [3]. One cools the system below T_{SG} and waits for a “waiting time” t_w . The system is then perturbed in some way, e.g. by applying a magnetic field, and the subsequent response is measured. It is found that the nature of the response depends on t_w , providing clear evidence that the system was not in equilibrium.

More complicated temperature protocols are possible, which have led to surprising results. For example, one can cool smoothly below T_{SG} and wait at a temperature T_1 , say, before cooling further, and then warming back up through T_{SG} this time without waiting at T_1 . While waiting at T_1 during the cooling process, the data shows a drift with time, and on warming, one finds a similar feature at T_1 even though the system did not wait there. This “memory” effect [4] is still not well understood, and neither is “rejuvenation”, the fact that aging at one temperature does not help equilibration at a lower temperature [4].

On the theoretical side, there is a mean field solution due to Parisi [5,6] which, following Sherrington and Kirkpatrick [7], is the exact solution of an EA-like model with infinite range interactions. One finds a finite spin glass transition temperature T_{SG} .

Most of what we know about short range short-range (EA) models in three dimensions has come from simulations on Ising systems, which also indicate a finite T_{SG} , as we will see below. However, less is known about vector spin glass models and these will be the main focus of the rest of the talk.

While the existence of a phase transition in three-dimensions is not in serious dispute, at least for Ising spins, the nature of the equilibrium state below T_{SG} has been much more controversial. An experimental system is not in equilibrium below T_{SG} , so we ultimately need a theory for non-equilibrium behavior. However, to do this, we presumably need to know the equilibrium state towards which the system is trying to reach but never does. Two main proposals have been made for the nature of the equilibrium spin glass state:

- “Replica Symmetry Breaking (RSB), which is like the Parisi [5,6] mean field solution, and
- The “droplet picture” (DP) of Fisher and Huse [8,9].

These differ in the nature of the large-scale, low-energy excitations, whose energy ΔE scales as

$$\Delta E \propto \ell^\theta, \tag{8}$$

where ℓ is the linear size of the excitation and θ is a “stiffness” exponent. RSB and DP have different predictions for θ :

- RSB, $\theta = 0$ for some excitations.
- DP, $\theta > 0$ (but small, around 0.2 for 3d Ising).

Since in both scenarios θ is zero or very small, a lot of cancellation occurs in the energy to flip a cluster of spins. Hence, a characteristic feature of spin glasses is that there are excitations which involve a large number of spins but which cost very little energy.

In three dimensions, numerics, which are inevitably on small lattice sizes, seem to fit best an intermediate (TNT) scenario [10,11] scenario. In two dimensions, where $\theta <$

0 and consequently $T_{SG} = 0$, larger sizes can be studied and it seems that the droplet theory works, though there are significant corrections to scaling for various quantities, see e.g. Refs. [12,13] for recent discussions.

A lot of attention in spin glasses has been spent on (i) the nature of the phase transition and (ii) the nature of the spin glass phase below T_{SG} . For both problems, most theory has been on Ising systems though the vector nature of the spins may be relevant. In the rest of this talk I will discuss the nature of the phase transition in *vector* spin glass models.

2. Vector Spin Glasses

Most theory has been done for the Ising ($S_i = \pm 1$) spin glass, where there is clear evidence for a finite T_{SG} . The best evidence is from finite size scaling (FSS) of the correlation length by Ballesteros et al. [14] This technique is discussed further below. However, many experimental systems, such as CuMn described above, are closer to an isotropic vector spin glass (in which \mathbf{S}_i is a vector), where the theoretical situation is less clear.

Old Monte Carlo simulations [15] found that T_{SG} , if it occurs at all, must be very low, and this was interpreted as being evidence for $T_{SG} = 0$. Motivated by this, Kawamura [16–19] argued that $T_{SG} = 0$ but there can be a glass-like transition at $T = T_{CG}$ in the “chiralities” (i.e. vortices). This implies *spin–chirality decoupling*. However, the possibility of finite T_{SG} has been raised by various authors, e.g. Maucourt and Gempel [20], Akino and Kosterlitz [21], Granato [22], Matsubara et al. [23,24], and Nakamura et al. [25]. Since the most successful approach for the Ising spin glass was the FSS scaling analysis of the correlation length, Lee and I decided to perform analogous calculations for vector spin glasses, investigating the correlation lengths of *both* the spins and chiralities.

There is an important difference between chiralities in frustrated and unfrustrated systems. In unfrustrated systems the ground state is collinear and so chirality needs to be *thermally excited*. Such thermally activated chiralities (vortices) are responsible for the Kosterlitz-Thouless-Berezinskii transition in the 2d XY ferromagnet. However, in spin glasses, chiralities are *quenched in* at low-T because the ground state is non-collinear as a result of the disorder and frustration.

To define chirality we follow Kawamura [17,18]:

$$\kappa_i^\mu = \begin{cases} \frac{1}{2\sqrt{2}} \sum_{\langle l,m \rangle} \text{sgn}(J_{lm}) \sin(\theta_l - \theta_m), & \text{XY } (\mu \perp \text{square}), \\ \mathbf{S}_{i+\hat{\mu}} \cdot \mathbf{S}_i \times \mathbf{S}_{i-\hat{\mu}}, & \text{Heisenberg,} \end{cases} \quad (9)$$

see Fig. 3, where for the XY case i refers to the plaquette indicated, and for the Heisenberg model, i refers to the middle of the three sites.

To determine the correlation lengths of the spins and chiralities we need to Fourier transform the appropriate correlation functions:

$$\begin{aligned} \chi_{SG}(\mathbf{k}) &= \frac{1}{N} \sum_{i,j} [\langle \mathbf{S}_i \cdot \mathbf{S}_j \rangle^2]_{\text{av}} e^{i\mathbf{k} \cdot (\mathbf{R}_i - \mathbf{R}_j)}, & (\text{spins}), \\ \chi_{CG}^\mu(\mathbf{k}) &= \frac{1}{N} \sum_{i,j} [\langle \kappa_i^\mu \kappa_j^\mu \rangle^2]_{\text{av}} e^{i\mathbf{k} \cdot (\mathbf{R}_i - \mathbf{R}_j)}, & (\text{chiralities}). \end{aligned} \quad (10)$$




Figure 3. An illustration of chirality for XY and Heisenberg spin glasses.

Note that $\chi_{nl} \sim \chi_{SG}(\mathbf{k} = 0)$, which is essentially the “correlation volume” of the spins.

We determine the spin glass correlation length of the finite-size system, ξ_L , from the Ornstein Zernicke equation:

$$\chi_{SG}(\mathbf{k}) = \frac{\chi_{SG}(\mathbf{0})}{1 + \xi_L^2 \mathbf{k}^2 + \dots}, \quad (11)$$

by fitting to $\mathbf{k} = 0$ and $\mathbf{k} = \mathbf{k}_{\min} = \frac{2\pi}{L}(1, 0, 0)$. The precise formula is

$$\xi_L = \frac{1}{2 \sin(k_{\min}/2)} \left(\frac{\chi_{SG}(\mathbf{0})}{\chi_{SG}(\mathbf{k}_{\min})} - 1 \right)^{1/2}. \quad (12)$$

The chiral glass correlation length of the system, $\xi_{c,L}^\mu$, is determined in an analogous way.

In order to locate the transition we use the technique of finite-size scaling (FSS). The basic assumption of FSS is that the size dependence comes from the ratio L/ξ_{bulk} where

$$\xi_{\text{bulk}} \sim (T - T_{SG})^{-\nu} \quad (13)$$

is the *bulk* correlation length. In particular, the *finite-size* correlation length is expected to vary as

$$\frac{\xi_L}{L} = X \left(L^{1/\nu} (T - T_{SG}) \right), \quad (14)$$

since ξ_L/L is dimensionless (and so has no power of L multiplying the scaling function X). Hence data for ξ_L/L for different sizes should intersect at T_{SG} and splay out below T_{SG} . Similarly, data for $\xi_{c,L}$ should intersect at T_{CG} .

3. Results

Figure 4 shows that this works well for the Ising spin glass. This data, which is for the spin glass correlation length divided by L , shows clear intersections and hence evidence




Figure 4. Data for the correlation length of the Ising spin glass showing clear evidence for a transition at $T_{SG} \simeq 1.00$.

for a transition, at $T_{SG} \simeq 1.00$. Furthermore, the data splay out again on the low- T side demonstrating that there is spin glass order below T_{SG} . This is data for the Gaussian distribution. The technique of determining T_{SG} by FSS of ξ_L was first used by Ballesteros et al. [14] who took the “ $\pm J$ ” distribution in which $J_{ij} = \pm 1$ with equal probability. This has a somewhat higher transition temperature, $T_{SG} \simeq 1.14$.

Prior to the work of Ballesteros et al., determination of T_{SG} generally used the “Binder ratio”, a dimensionless ratio of the moments of the order parameter distribution which has a finite size scaling of the same form as in Eq. (14). However, this gives much less convincing demonstration of a transition, see Fig. 5 which shows data from Marinari et al. [26] for the Gaussian distribution.

We have seen that the best method for studying the transition in the Ising spin glass is FSS of the correlation length. We now apply this to the spin glass with vector spins. Similar results were obtained [27] for both the XY and Heisenberg models. Here, for conciseness, we just present results for the Heisenberg case.

Figure 6 shows data for ξ_L/L . It has some additional data beyond that given in Lee and Young [27]. The data intersects and splays out again at low temperatures indicating a finite-temperature spin glass transition. The inset shows that the data can be collapsed reasonably according to the FSS form in Eq. (14) with $T_{SG} \simeq 0.16$, $\nu \simeq 1.2$.

Figure 7 shows data for the chiral correlation length. There are actually two such lengths depending upon whether the wavevector \mathbf{k}_{\min} in Eq. (12) is parallel or perpendicular to the line of spins shown in Fig. 3 for the Heisenberg case. The main figure in Fig. 7 shows the parallel correlation length and the inset shows the perpendicular correlation length. Apart from the smallest size, the data intersect pretty well. Furthermore, the transition temperature




Figure 5. Data for the Binder ratio length of the Ising spin glass with Gaussian interactions, from Marinari et al. [26]. The data merge but do not splay out strongly on the low- T side, unlike the results for the correlation length shown in Fig. 4.




Figure 6. Data for the spin glass correlation length of the Heisenberg spin glass divided by L (based on Ref. [27] with some additional data).




Figure 7. Data for the chiral glass correlation length of the Heisenberg spin glass divided by L (based on Ref. [27] with some additional data).

T_{CG} seems to be equal to T_{SG} , namely about 0.16.

We conclude that a *direct* study of the correlation lengths indicates that there is a single phase transition at which both spins and chiralities order in vector spin glasses.

4. Conclusions

It is interesting to see how the spin glass transition temperature varies with the number of spin components m . To compare different values of m it is necessary to note that there is an m dependence for T_{SG} even in mean field theory: $T_{SG}^{MF} = \sqrt{z}/m$ where z ($= 6$ here) is the number of neighbors. Hence we show below values for T_{SG}/T_{SG}^{MF} determined from the numerics:

m	model	T_{SG}^{MF}	T_{SG}	T_{SG}/T_{SG}^{MF}
1	(Ising)	2.45	1.00	0.41
2	(XY)	1.22	0.34	0.28
3	(Heisenberg)	0.82	0.16	0.20

We see that T_{SG}/T_{SG}^{MF} is small and decreases further with increasing m . Physically, this means that fluctuation effects are large and get larger with increasing m . The data suggest that perhaps $T_{SG} = 0$ for $m = \infty$. Preliminary results by Dhar, Lee and the author [28], indicate that this is the case.

To conclude, I believe that one important question, whether or not an isotropic Heisenberg spin glass has a finite temperature spin glass transition, has been answered in the affirmative. However, the nature of the putative equilibrium state below T_{SG} , towards which the system evolves but never reaches, as well as non-equilibrium phenomena such as aging and rejuvenation, remain to be fully understood.

ACKNOWLEDGMENTS

This work was done in collaboration with Lik Wee Lee and I would like to thank him for many valuable discussions. I acknowledge support from the National Science Foundation under grant DMR 0337049.

References

- [1] S. F. Edwards and P. W. Anderson, *J. Phys. F*, 5, 965 (1975).
- [2] R. Omari, J. J. Prejean, and J. Souletie, *J. de Physique*, 44, 1069 (1983).
- [3] P. Nordblad and P. Svendlidh, in *Spin glasses and random fields*, edited by A. P. Young, (World Scientific, Singapore, 1998).
- [4] K. Jonason, E. Vincent, J. Hammann, J. Bouchaud, and P. Nordblad, *Phys. Rev. Lett.* 81, 3243 (1998).
- [5] G. Parisi, *J. Phys. A*, 13, 1101 (1980).
- [6] G. Parisi, *Phys. Rev. Lett.* 50, 1946 (1983).
- [7] D. Sherrington and S. Kirkpatrick, *Phys. Rev. Lett.* 35, 1972 (1975).
- [8] D. S. Fisher and D. A. Huse, *Phys. Rev. Lett.* 56, 1601 (1986).
- [9] D. S. Fisher and D. A. Huse, *Phys. Rev. B* 38, 386 (1988).
- [10] F. Krzakala and O. C. Martin, *Phys. Rev. Lett.*, 85, 3013 (2000).
- [11] M. Palassini and A. P. Young, *Phys. Rev. Lett.*, 85, 3017 (2000).
- [12] A. K. Hartmann and M. A. Moore, *Phys. Rev. Lett.*, 90, 127201 (2003).
- [13] H. G. Katzgraber, L. W. Lee and A. P. Young, (cond-mat/0402037).
- [14] H. G. Ballesteros, A. Cruz, L. A. Fernandez, V. Martin-Mayor, J. Pech, J. J. Ruiz-Lorenzo, A. Tarancon, P. Tellez, C. L. Ullod, and C. Ungil, *Phys. Rev. B* 62, 14237 (2000).
- [15] S. Jain and A. P. Young, *J. Phys. C* 19, 3913 (1986).
- [16] H. Kawamura and M. Tanemura, *Phys. Rev. B*, 36, 7177 (1987).
- [17] H. Kawamura and M. S. Li, *Phys. Rev. Lett.* 87, 187204 (2001).
- [18] H. Kawamura, *Phys. Rev. Lett.* 80, 5421 (1998).
- [19] K. Hukushima and H. Kawamura, *Phys. Rev. E* 61, R1008 (2000).
- [20] J. Maucourt and D. R. Grempel, *Phys. Rev. Lett.* 80, 770 (1998).
- [21] N. Akino and J. M. Kosterlitz, *Phys. Rev. B* 66, 054536 (2002).
- [22] E. Granato, *J. Magn. Magn. Matter.* 226, 366 (2000).
- [23] F. Matsubara, T. Shirakura, and S. Endoh, *Phys. Rev. B* 64, 092412 (2001).
- [24] S. Endoh, F. Matsubara, and T. Shirakura, *J. Phys. Soc. Jpn.* 70, 1543 (2001).
- [25] T. Nakamura and S. Endoh, *J. Phys. Soc. Jpn.* 71, 2113 (2002).
- [26] E. Marinari, G. Parisi, and J. J. Ruiz-Lorenzo, *Phys. Rev. B* 58, 14852 (1998).
- [27] L. W. Lee and A. P. Young, *Phys. Rev. Lett.* 90, 227203 (2003).
- [28] A. Dhar, L. W. Lee and A. P. Young, (unpublished).